

Appl. No. 10/712,967
Amdt dated October , 2006
Reply to Office Action of May 26, 2006
Att. Docket No.: 1279-291N1

Filing date: November 14, 2003
Applicant Name: Timothy J. Deming
Examiner: David Lukton
Art Unit: 1654

RECEIVED
CENTRAL FAX CENTER

OCT 25 2006


Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of the claims in the application:

Listing of Claims:

Claims 1-15 (canceled)

Claim 16 (currently amended) A method of polymerizing an aminoacid-N-carboxyanhydride monomer[[s]] ~~having a ring with a O-C₅ and a O-C₂ anhydride bond comprising: wherein the N-carboxyanhydride monomer comprises the following ring:~~



said method comprising:

- (a) combining a first NCA monomer with an initiator molecule complex comprised of:
 - (i) a low valent metal capable of undergoing an oxidative addition reaction wherein the oxidative addition reaction formally increases the oxidation state of the low valent metal by two electrons; and
 - (ii) an electron donor comprising a Lewis base;
- (b) allowing the initiator molecule complex to:

Appl. No. 10/712,967
 Amdt dated October , 2006
 Reply to Office Action of May 26, 2006
 Att. Docket No.: 1279-291NI

Filing date: November 14, 2003
 Applicant Name: Timothy J. Deming
 Examiner: David Lukton
 Art Unit: 1634

(i) open the ring of the first NCA through oxidative addition across either the O-C₅ or O-C₂ anhydride bond;

(ii) combine with a second NCA monomer, to form ~~an~~ a polyaminoacid chain containing a first amido containing metallacycle; and


(c) allowing a third NCA monomer to combine with the first amido containing ~~metallacycle~~ metallacycle so that:

(i) the amido nitrogen of the first amido containing ~~metallacycle~~ metallacycle attacks the carbonyl carbon of the NCA and the third NCA is added to the polyaminoacid chain; and

(ii) ~~the a second~~ a second amido containing ~~metallacycle~~ metallacycle is ~~regenerated~~ generated.

Claim 17 (currently amended) A method of adding an aminoacid-N-carboxyanhydride to a polyaminoacid chain having an amido containing metallacycle end group wherein the metallacycle end group is of the general formula:

~~The method of claim 16 wherein the amido-containing metallacycle is of the general formula:~~



wherein M is the low valent transition metal;

L is the Lewis Base ligand;

Appl. No. 10/712,967
Amdt dated October , 2006
Reply to Office Action of May 26, 2006
Att. Docket No.: 1279-291N1

Filing date: November 14, 2003
Applicant Name: Timothy J. Deming
Examiner: David Lukton
Art Unit: 1654

R1, R2 and R3 comprises a side chain of an amino acid selected from the group consisting of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine; and

R4 is the polyaminoacid chain[(.)], and

n is an integer from 1 to 4.

Claim 18 (original) The method of claim 16 wherein the efficiency of the initiator is controlled by allowing the reaction to proceed in a solvent selected for its ability to influence the reaction.

Claim 19 (original) The method of claim 17 wherein the solvent is selected from the group consisting of ethyl acetate, toluene, dioxane, acetonitrile, THF and DMF.

Claim 20 (original) The method of claim 16, wherein the α -aminoacid-N-carboxyanhydride monomer is selected from the group consisting of alanine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine and valine.

Claims 21-62 (canceled)